

# RSC Mechanochemistry

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: H. Chen, Z. Guo, D. Feng, X. Jin and F. Guo, *RSC Mechanochem.*, 2025, DOI: 10.1039/D5MR00057B.



This is an Accepted Manuscript, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about Accepted Manuscripts in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this Accepted Manuscript or any consequences arising from the use of any information it contains.

# Mechanochemical One-Pot Synthesis and Solid-State Transformation of Cobalt(II) Schiff Base Complexes: A Green Route to Tailored Coordination Architectures

Hongguang Chen, Zhenwei Guo, Daming Feng\*, Xudong Jin\*, Fang Guo\*

College of Chemistry, Liaoning University, Shenyang 110036, China.

E-mail: dmfeng@lnu.edu.cn; [jinxudong@lnu.edu.cn](mailto:jinxudong@lnu.edu.cn); [fguo@lnu.edu.cn](mailto:fguo@lnu.edu.cn).

## Abstract

Schiff base complexes, traditionally synthesized via time-consuming, solvent-intensive solution methods, are pivotal in coordination chemistry but face limitations in accessing diverse architectures and sustainable scalability. Mechanochemistry has emerged as a solvent-free alternative, yet its potential to drive multicomponent reactions with precise control over metal-ligand coordination modes remains underexplored. Herein, we propose a mechanochemical one-pot synthesis strategy that synergistically integrates condensation, metal coordination, and deprotonation-dehalogenation reactions to fabricate Schiff base Co(II) complexes. Utilizing adamantylamine, 5-halosalicylaldehyde, and  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  as precursors, mechanical forces drive the self-ordering of building blocks, enabling ultrahigh selective coordination and complex chemical processes. The approach efficiently yielded 12 Co(II) complexes, including the  $\kappa^1\text{-O}$ -monodentate  $\text{CoCl}_2(\text{HL})_2$  and  $\kappa^2\text{-O}$ , N-bidentate  $\text{CoL}_2$ , which are typically challenging to access via conventional solution methods. Remarkably, the reactions achieved full conversion within 10 minutes, underscoring the rapidity and sustainability of mechanochemistry. Mechanical activation unlocked dormant reactivity in reactants, facilitating pathways otherwise inaccessible in solution. Furthermore, reversible solid-state transformations between complexes were demonstrated through dehydrohalogenation-hydrohalogenation processes. Specifically,  $\text{CoCl}_2(\text{HL})_2$  converts to  $\text{CoL}_2$  via cleavage of N–H and Co–Cl bonds and subsequent Co–N bond formation, while  $\text{CoL}_2$



reverts upon HCl absorption during grinding. This work highlights the utility of mechanochemistry in simplifying synthetic procedures, enhancing reaction complexity, and enabling green, solvent-free synthesis. By elucidating pathways for one-pot synthesis and solid-state transformations, it establishes mechanochemistry as a versatile and sustainable route for designing advanced coordination complexes with tailored architectures.

**KEYWORDS:** Mechanochemistry; One-pot synthesis; Schiff base complexes; Co(II) coordination chemistry; Green synthesis.

## Introduction

The utilization of mechanical energy to induce or promote chemical transformations for driving chemical reactions has emerged as a significant approach in chemical synthesis.<sup>1</sup> This method notably reduces, or even entirely eliminates, the reliance on solvents, offering a more sustainable alternative to traditional solvent-based synthesis, and is progressively being recognized as an innovative green synthesis technique.<sup>2, 3</sup> As a pivotal aspect of mechanochemical synthesis, grinding methods can effectively facilitate chemical reactions under solvent-free (NG) or liquid-assisted grinding (LAG) conditions.<sup>4</sup> These methods are particularly advantageous for reactions involving substances with poor solubility or substantial discrepancies in solubility and have garnered considerable attention in the field of complex organic ligand preparation and their metal complexes.<sup>5</sup> Furthermore, grinding methods can alter established chemical reaction pathways and selectivity (yielding reaction pathways distinct from the thermodynamic products in solution),<sup>6</sup> thereby overcoming the constraints of traditional solution synthesis and allowing the formation of product structures that are challenging to generate or thermodynamically unfavorable in conventional solution methods, thus providing a potent tool for the discovery of new chemical reactivities, complex configurations, and transformation pathways.<sup>7</sup>


Schiff bases, characterized by their unique electronic properties, are capable of coordinating with various metal ions to form complexes with diverse geometries and stereoselectivities<sup>8</sup>, an ability that positions them prominently in the field of coordination chemistry and demonstrates significant advantages in various fields<sup>9</sup>, including biomedicine<sup>10</sup>, photo-oxidation catalysis<sup>11</sup>, and chemical sensing.<sup>12</sup> However, traditional solution synthesis methods face numerous challenges in the preparation of Schiff bases and their corresponding complexes. Variations in the solubility of



the reactants often require harsh conditions at high temperatures via thermodynamic control issues and low yields.<sup>13</sup> Furthermore, the imine bond is prone to hydrolysis or oxidation, which compromises the stability of Schiff bases, particularly in monodentate Schiff base complexes.<sup>14</sup> Mechanochemistry now facilitates the solvent-free synthesis of chelate complexes containing C=N ligands via mechanically driven imine condensation combined with in-situ coordination, providing a rapid and environmentally friendly pathway to Schiff-base complexes.<sup>15, 16,17</sup> Consequently, a significant area of research has developed, concentrating on the creation of novel Schiff-base complexes characterized by environmentally sustainable, controllable, and diverse structures, while also exploring the reversible solid-state conversion of their coordination configurations and their potential applications in biomedicine and other fields.<sup>18</sup> Nevertheless, a significant barrier to the advancement of structurally diverse coordination systems is the absence of strategies to concurrently manage multiple interrelated dynamic processes across various architectural factors, including reversible covalent bond formation, reversible hydrogen dehalogenation, and dynamic metal-ligand coordination.<sup>19, 20</sup> Previous studies have documented hydrogen dehalogenation methods and the resultant changes in ligand structure before and after the reaction (Scheme 1a).<sup>21</sup> Notably, many Schiff base structures exhibit intramolecular bonding<sup>22, 23</sup> (Figure S1), suggesting that dehydrohalogenation reactions may occur when reactants are appropriately oriented, potentially leading to the development of novel Schiff base complexes through structural transformations<sup>24</sup> (Scheme 1b).

In this context, it is essential to explore facile and atom-economical multicomponent reaction strategies that simultaneously modulate the dynamic interactions between hydrogen bonding sites, imine covalent sites, and coordination sites through mechanochemical one-pot methods, which aims to promote self-ordering behaviors for the controllable preparation of Schiff complexes with various coordination modes. In particular, the construction of Schiff base bidentate and monodentate complex molecules exhibiting antimicrobial properties can be achieved using adamantyl compounds (e.g., adamantanamine, adamantylethylamine, and memantine), which are employed in the treatment of influenza A virus, Parkinson's disease, and for the prevention of early influenza infections, as well as the treatment of severe Alzheimer's disease.<sup>25, 26</sup> Furthermore, controlling the presence or absence of intracellular hydrogen bonding N-H-O is crucial for modulating cobalt (II) metal coordination modes in a controllable manner.





**Scheme 1.** Hydrodehalogenation methods have been reported (a). Mechanochemical one-pot process for the synthesis of  $\text{CoCl}_2(\text{HL})_2$  and  $\text{CoL}_2$ , along with a reversible hydrogen dehalogenation scheme (b).

This study demonstrates that a series of monodentate Co(II) complexes, designated as  $\text{CoCl}_2(\text{HL})_2$  ( $L = 2-((\text{adamantan-1-yl})\text{-imino-methyl})\text{-4-chlorophenol}$  and five additional ligand derivatives), can be synthesized through either sequential or one-pot mechanochemical synthesis *via* neat grinding of precursors A (amantadine, memantine, rimantadine), B (5-chlorosalicylaldehyde, 5-bromosalicylaldehyde) and  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  in 2:2:1 molar ratio. These complexes are obtained quantitatively within 5 minutes under solvent-free conditions, as depicted in Scheme 1. By incorporating NaOH into the same reactant mixture and extending the grinding time to 10 minutes, a series of  $\kappa^2\text{-O, N}$ -bidentate  $\text{CoL}_2$  complexes can also be prepared using the same mechanochemical protocol. Mechanochemical dehydrochlorination of  $\kappa^1\text{-O}$ -monodentate  $\text{CoCl}_2(\text{HL})_2$  enables the synthesis of  $\kappa^2\text{-O, N}$ -bidentate  $\text{CoL}_2$  complexes, while exposure to HCl allows reversible transformation of  $\text{CoL}_2$  back to  $\kappa^1\text{-O}$ -monodentate  $\text{CoCl}_2(\text{HL})_2$ . Both transformations yield high purity products within 5 minutes. Notably,  $\kappa^1\text{-O}$ -monodentate  $\text{CoCl}_2(\text{HL})_2$  complexes are challenging to synthesize and unstable in the solution, necessitating a




solvent-free approach. This approach thus provides a novel pathway for preparing solvent-sensitive monodentate complexes.

## Results and Discussion

Six  $\kappa^1$ -O-monodentate  $\text{CoCl}_2(\text{HL})_2$  complexes were synthesized through a one-pot mechanochemical process by mixing and grinding component A (adamantanamine, adamantylethylamine, and mecamantanamine), component B (5-chlorosalicylaldehyde and 5-bromosalicylaldehyde), and component C ( $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ ) in a molar ratio of 2:2:1 over a duration of 10 minutes, resulting in the green powder product. When ground under the same reaction conditions and utilizing the same reactant system with the addition of 2 equivalents of NaOH resulted in the formation of six distinct  $\kappa^2$ -O, N-bidentate complexes  $\text{CoL}_2$ , characterized by a red powder product which differed from the previously observed  $\kappa^1$ -O-monodentate complexes, were formed alongside NaCl, as confirmed by the presence of entirely new diffraction peaks in the powder X-ray diffraction (PXRD) pattern shown in Figure 1 and Figure S2 and S3. The analysis of the PXRD patterns of the synthesized products revealed that the diffraction peak positions and intensities of the ground products closely matched those of the crystal simulations and the Schiff base cobalt complex reported in the literature. This strongly confirms the success of the mechanochemical multicomponent one-pot method for synthesizing the metal-organic complexes, thereby demonstrating the feasibility and validity of this synthesis strategy within this system. The PXRD patterns of the remaining five pairs of complexes are presented in Figures S4-8. FT-IR analysis reveals that hydrogen-bond-mediated  $\kappa^1$ -O-monodentate  $\text{CoCl}_2(\text{HL})_2$  exhibits a loss of O–H at  $3430\text{ cm}^{-1}$  and N–H at  $3220\text{ cm}^{-1}$ , in contrast to the dehydrochlorination process yielding  $\kappa^2$ -O, N-bidentate  $\text{CoL}_2$ , which shows a red-shifted C=N at  $1600\text{ cm}^{-1}$  and a new Co–N bond at  $480\text{ cm}^{-1}$  (Figures S25-S36). High-resolution mass spectrometry (HRMS-ESI) tests indicate that the experimental values are consistent with their theoretical values, demonstrating that the structure of the complexes synthesized via mechanochemical one-pot methods corresponds to the structures obtained through quantitative synthesis. In summary, mechanochemistry can efficiently and controllably synthesize  $\kappa^1$ -O-mono- and  $\kappa^2$ -O, N-bidentate cobalt Schiff base complexes.








**Figure 1.** PXRD patterns of  $\kappa^1$ -O-monodentate **CoCl<sub>2</sub>(HL<sub>2</sub>)<sub>2</sub>** complexes and  $\kappa^2$ -O, N-bidentate complexes **CoL<sub>2</sub>** synthesized by mechanochemical one-pot process and dehalogenation of hydrogen.

Among all possible combinations of initial components, if further condensation of an imine-containing ligand can provide an optimal coordination environment for the metal, the formation of this ligand can be selectively facilitated by its preferential binding to the transition metal, allowing for the formation of complexes with different configurations selectively.<sup>27</sup> During the one-pot synthesis, the condensation covalent reaction of ammonia and aldehyde, the metal coordination reaction, and the deprotonated dehalogenation reaction were placed in the same reaction system to act synergistically, and underwent synergistic and complex chemical processes facilitated by mechanical force, inducing self-classification and ultra-high-selective coordination, and successfully realizing the highly efficient construction of cobalt metal-organic complexes. The mechanochemical synthesis of Schiff base ligands and their cobalt  $\kappa^1$ -O-mono- and  $\kappa^2$ -O, N-bidentate complexes can be fully converted after 10 minutes, and this study highlights the feasibility of mechanochemistry in the sustainable synthesis of complex organic ligands and their




metal complexes. Compared to the solution method, the mechanochemical one-pot synthesis has advantages such as solvent-free conditions, rapid processing, minimal energy consumption, and by-products consisting solely of NaCl, as detailed in Table S5. More importantly, the UV spectra of the monodentate complexes in six common organic solvents reveal that, in addition to the peak at 325 nm, a new red-shifted absorption peak appears at 390 nm in organic protic solvents, compared to the characteristic absorption peaks in non-protic solvents (Figures S37-S42). This indicates a change in the coordination mode of the complex in protic solvents, suggesting rapid dehydrochlorination and coordination rearrangement. Therefore, mechanochemical methods can synthesize monodentate cobalt complexes that are not achievable through conventional solution methods.

The above grinding product can be readily crystallized under appropriate solvents, giving rise to crystals for X-ray single crystal structural determination. As illustrated in Table S1 and S2, the formation of the two lattices is contingent upon the adamantamine/metamantamine and adamantane ethylamine ligands, where the presence or absence of methylene groups in proximity to the reaction site of the amino molecule is crucial for the lattice formation. The crystals of the methylene-free  $\kappa^1$ -O-monodentate complexes formed by the Schiff base ligands **L1-L4**, derived from amantadine and mecamtadine, are labeled as **CoCl<sub>2</sub>(HL1)<sub>2</sub>·CH<sub>3</sub>CN**, **CoCl<sub>2</sub>(HL2)<sub>2</sub>·CH<sub>3</sub>CN**, **CoCl<sub>2</sub>(HL3)<sub>2</sub>**, and **CoCl<sub>2</sub>(HL4)<sub>2</sub>**, respectively, exhibit isostructures belonging to the space group *P2<sub>1</sub>/c* (see Table S1). Notably, powder X-ray diffraction patterns reveal significant differences between **CoCl<sub>2</sub>(HL1)<sub>2</sub>·CH<sub>3</sub>CN**, **CoCl<sub>2</sub>(HL2)<sub>2</sub>·CH<sub>3</sub>CN** and **CoCl<sub>2</sub>(HL3)<sub>2</sub>**, **CoCl<sub>2</sub>(HL4)<sub>2</sub>**, indicating that structural rearrangements occurred with solvent participation (Figure 2 and Figure S9). Conversely, the adamantylethylamine  $\kappa^1$ -O-monodentate complexes **CoCl<sub>2</sub>(HL5)<sub>2</sub>** and **CoCl<sub>2</sub>(HL6)<sub>2</sub>**, which contain methylene groups, share the same space group *C2/c*.








**Figure 2.** PXRD spectra of crystals of  $\text{CoCl}_2(\text{HL1})_2 \cdot \text{CH}_3\text{CN}$ ,  $\text{CoCl}_2(\text{HL2})_2 \cdot \text{CH}_3\text{CN}$ ,  $\text{CoCl}_2(\text{HL3})_2$  and  $\text{CoCl}_2(\text{HL4})_2$  obtained by recrystallisation.

Herein, we utilize  $\text{CoCl}_2(\text{HL2})_2 \cdot \text{CH}_3\text{CN}$  as a representative example to describe the crystal structures of  $\kappa^1$ -O-monodentate Schiff base complexes formed by ligands **L1-L4**. The crystal crystallizes in the monoclinic  $P2_1/c$  space group, with one asymmetric unit consisting of a  $\text{CoCl}_2(\text{HL2})_2$  and one acetonitrile molecule. As seen in Figure 3, the cobalt ion coordinates with two singly adjacent protonated HL molecules and two chlorine ions simultaneously, forming a four-coordinated configuration with an irregular tetrahedral spatial geometry ( $\tau_4 = 0.8998$ ), where each HL acts as a  $\kappa^1$ -O-monodentate coordination site via its phenolic oxygen atoms. The corresponding Co-O and Co-Cl bond lengths are 1.9610 Å (Co-O1) and 1.9530 Å (Co-O2), 2.2280 Å (Co-Cl1) and 2.2807 Å (Co-Cl2), the two hydroxyl groups form coordination dihedral angles of 81.26°. The C=N bond keeps the planarity with bromosalicylaldehyde ring, with the dihedral angle of 8.00° and 3.09°. The protonated hydroxyl forms intramolecular N-H $\cdots$ O hydrogen bonds (1.982 Å, 1.900 Å) with the N of imine, which is commonly existed in most Schiff ligand. The  $\text{CoCl}_2(\text{HL})_2$  units are connected with each other through extensive C-H $\cdots$ Cl and C-H $\cdots$ Br hydrogen bonding, and the included acetonitrile molecule is linked with  $\text{CoCl}_2(\text{HL})_2$  unit via C-H $\cdots$ N and C-H $\cdots$ Cl weak interactions to keep the stability of the structure (Figure S10). The




coordination and stacking patterns of  $\text{CoCl}_2(\text{HL1})_2 \cdot \text{CH}_3\text{CN}$  and  $\text{CoCl}_2(\text{HL2})_2 \cdot \text{CH}_3\text{CN}$  are identical (Figure S11). Both  $\text{CoCl}_2(\text{HL3})_2$  and  $\text{CoCl}_2(\text{HL4})_2$  exhibit identical coordination patterns and involve robust intramolecular hydrogen bonds between the imino nitrogen and the phenolic oxygen, resulting in the formation of an additional six-membered ring (Figure S12). The structures are more tightly stacked as the asymmetric unit does not contain solvent molecules.



**Figure 3.** Crystal structure and their packing for  $\text{CoCl}_2(\text{HL2})_2$

The  $\kappa^1\text{-O}$ -monodentate complex crystals formed by the adamantylethane Schiff base ligands **L5** and **L6** are labeled as  $\text{CoCl}_2(\text{HL5})_2$  and  $\text{CoCl}_2(\text{HL6})_2$ , respectively. These complexes are isostructural, sharing the same space group  $C2/c$ , with each asymmetric unit containing half of a complex molecule and exhibiting similar unit cell parameters, as shown in Table S2. In the crystal, the cobalt ions form a four-coordination configuration characterized by irregular tetrahedra ( $\tau_4 = 0.89$ ), involving two individual protonated HL molecules and two chloride ions Figure 4. Notably, the two  $\text{C}=\text{N}$  bonds remain approximately planar with bromosalicylaldehyde; however, the distance between the two nitrogen atoms is 8.02 Å. The two hydroxyl groups create coordination dihedral angles of 69.09° and 80.98°. Furthermore, two intramolecular  $\text{N-H}\cdots\text{O}$  hydrogen bonds are formed, with bond lengths of 1.948 Å and 1.955 Å. The neighboring complex molecules are primarily interconnected through  $\text{C-H}\cdots\text{Cl}$  weak interactions, resulting in chain stacking along the C-axis direction, which facilitates the maintenance of a relatively stable position and orientation within the crystal (Figure S13).






**Figure 4.** Crystal structure and their packing for  $\text{CoCl}_2(\text{HL5})_2$

The mechanochemical formed product from dehydrochlorination was further recrystallized from solvents, giving rise to the crystal's suitable crystals for single crystal X-ray diffraction, which was labelled as  $\text{CoL}_2$  ( $L = \text{L1-L6}$ ), respectively. As illustrated in Table S3, the presence or absence of methylene in the neighbourhood of the amino covalent reaction site also influences the lattice formation of  $\kappa^2\text{-O, N}$ -bidentate complexes. The methylene-free  $\kappa^2\text{-O, N}$ -bidentate complex crystals formed by ligands **L1**, **L3** and **L4**, show an isomorphic structure belonging to the  $Pbca$  space group. In contrast,  $\text{CoL2}_2 \cdot \text{DMSO}$ , which includes the solvent molecule DMSO, belongs to the  $P\bar{1}$  space group. Conversely, the  $\kappa^2\text{-O, N}$ -bidentate complexes  $\text{CoL5}_2$  and  $\text{CoL6}_2$ , derived from the methylene-containing ligands **L5** and **L6**, share the same  $P2_1/c$  space group. It is noteworthy that  $\text{CoL2}_2 \cdot \text{DMSO}$  has the same space group as the previously reported  $\text{CoL2}_2 \cdot \text{EtOH}$ , although with different cell parameters, it is considered to be a pseudo-polycrystalline form of  $\text{CoL2}_2$ .<sup>28</sup> The DMSO solvent molecules are released upon heating (Figure. S14) and can also be recovered through LAG grinding, and PXRD diffraction (Figure. S15) confirms the significant changes in the lattice structure.






**Figure 5.** Crystal structure and their packing for **CoL32·DMSO**.

Here we describe the crystal structures of  $\kappa^2$ -O, N-bidentate Schiff base complexes formed by the ligands L1-L4 using **CoL22·DMSO** as a representative. **CoL22·DMSO** crystallizes in the triclinic *P* $\bar{1}$  space group, with one asymmetric unit consisting of one **CoL22** and one dimethyl sulfoxide molecule, as seen in Figure 5. Although the cobalt ion maintains a four-coordination structure with an irregular tetrahedral spatial geometry ( $\tau_4 = 0.8500$ ), the Co ion coordinates with phenolic O atoms (O1 and O2) and imine N atoms (N1 and N2) of two singly deprotonated ligands simultaneously to form two nearly planar six-membered rings. The corresponding bond lengths of Co-N and Co-O are 1.9930 Å (Co-N1) and 2.0030 Å (Co-N2), 1.9606 Å (Co-O1) and 1.9130 Å (Co-O2). There are two types O-Co-N bond angles. Some are smaller than 109.5° for an ideal tetrahedron, while the others are wider. All bond angles of O-Co-O (112.90°) and N-Co-N (123.14°) are larger than 109.5°. The dihedral angles between the two chelate planes formed by Co-N-C-C-C-O are 86.76°, indicating that the two chelate planes are almost perpendicular. The **CoL22** unit is connected with each other through extensive C-H $\cdots$ Br hydrogen bonding. The included DMSO molecule is connected with CoL22 unit via C-H $\cdots$ O weak interactions to keep the stability of the structure (Figure S16). **CoL12**, **CoL32**, and **CoL42** exhibit the same cobalt bonding modes and coordination configurations as **CoL22·DMSO**, with the exception of the solvent-free molecules (Figure S17). All  $\kappa^2$ -O, N-bidentate coordination mode of the complexes does not involve intramolecular hydrogen bonds






because the ligand has been deprotonated, but there also exist two six-membered rings through a cobalt ion, two oxygen atoms and two nitrogen atoms.



**Figure 6.** Crystal structure and their packing for **CoL5<sub>2</sub>**.

The  $\kappa^2$ -O, N-bidentate complexes **CoL5<sub>2</sub>** and **CoL6<sub>2</sub>** crystallize in the monoclinic system, belonging to the  $P2_1/c$  space group, each asymmetric unit comprises one cobalt (II) complex. The central cobalt (II) in both **CoL5<sub>2</sub>** and **CoL6<sub>2</sub>** lies on a twofold rotation axis and is bonded to the oxygen and nitrogen donors of the two  $\kappa^2$ -O, N-bidentate ligands in a trans arrangement (Figure 6 and S18).<sup>29</sup> The complexes exhibit a distorted tetrahedral geometry ( $\tau_4 = 0.83$ ), as the  $\tau_4$  parameter is less than 0.85, indicating a configuration that is more distorted than that of other  $\kappa^2$ -O, N-bidentate complexes. The dihedral angle between the two coordination planes defined by O1–Co1–N1 and O2–Co1–N2 measures  $86.01^\circ$  for **CoL5<sub>2</sub>** and  $85.76^\circ$  for **CoL6<sub>2</sub>**. Additionally, the relatively large distance between the adamantylethylamine cages of the two ligands leads to an antcoordination phenomenon, where the imine-coordinated form predominates. The  $\kappa^2$ -O, N-bidentate coordination inhibits the formation of intramolecular hydrogen bonds due to the deprotonated ligands; however, two six-membered rings are also present, formed through a cobalt ion, two oxygen atoms, and two nitrogen atoms in both **CoL5<sub>2</sub>** and **CoL6<sub>2</sub>**.

The crystal structure reveals that the conversion of  $\kappa^1$ -O-monodentate complexes to  $\kappa^2$ -O, N-bidentate complexes occurs through the dehydrochlorination, accompanied by a reduction in the distance between the metal and the ligand. This process facilitates the ‘squeezing’ of ligands,



resulting in a more tightly packed  $\kappa^2$ -O, N-bidentate coordination configuration. The inherent instability of  $\kappa^1$ -O-monodentate molecules, which are loosely packed due to longer coordination bonds compared to  $\kappa^2$ -O, N-bidentate complexes, is further exacerbated by dehydrogenation stemming from the disruption of intramolecular hydrogen bonds. This dehydrogenation serves as the primary driving force for the transition from  $\kappa^1$ -O-monodentate to  $\kappa^2$ -O, N-bidentate coordination. Even in the absence of alkaline substances, hydrogen bonding can still occur in protonic solvents such as methanol and ethanol, which contain both hydrogen bond donors and acceptors. This interaction disrupts the N-H $\cdots$ O intramolecular hydrogen bonding, resulting in a rearrangement of the coordination mode that diminishes the selectivity for  $\kappa^1$ -O-monodentate complexes, ultimately leading to the exclusive formation of  $\kappa^2$ -O, N-bidentate complexes. Moreover, during this rearrangement process, the metal cation does not form a coordination bond with solvent molecules; instead, it acts as a strong hydrogen bond acceptor, facilitating the removal of the halogen anion and stabilizing the coordination rearrangement. The  $\kappa^1$ -O-monodentate complexes yielded single crystals with  $\kappa^2$ -O, N-bidentate coordination, except when recrystallized in acetonitrile solvents, thereby validating our design principle. In contrast, conventional solvothermal synthesis typically incorporates nitrogen and oxygen into their coordination spheres, resulting in the formation of only a single  $\kappa^2$ -O, N-bidentate coordination.<sup>30, 31</sup> This paper presents a solvent-free one-pot mechanochemical synthesis that mitigates the influence of solvents, thereby accelerating the formation of  $\kappa^1$ -O-monodentate coordination and preventing its rearrangement, allowing the synthesis of  $\kappa^1$ -O-monodentate complexes that cannot be achieved by solvent methods. Additionally,  $\kappa^2$ -O, N-bidentate complexes can be further synthesized through alkali grinding and dehalogenation of hydrogen, and the selective preparation of  $\kappa^1$ -O-monodentate and  $\kappa^2$ -O, N-bidentate coordination structures is facilitated by utilizing intramolecular hydrogen bonding to regulate the self-sorting process of mono- and  $\kappa^2$ -O, N-bidentate coordination.

Factors such as the denticity, electronic properties and steric site resistance of the ligand can modulate its binding strength to transition metals, which results from the unique electronic configuration of the ligand that imparts a specific coordination tendency to the transition metal.<sup>19, 32</sup> When the organic initial component, adamantanamine, is replaced by the more electron-donating adamantylethylamine and mecamtadine, or when the 4-substituent of salicylaldehyde is altered






from chlorine to bromine, the resulting  $\kappa^1$ -O-monodentate and corresponding  $\kappa^2$ -O, N-bidentate complexes do not exhibit significant changes in the coordination conformations of cobalt and the ligand, suggesting that remote electronic interactions do not substantially affect the binding between the metal and the ligand site. However, from the perspective of crystal stacking, the volume of the amino group significantly influences the crystal arrangement. The larger volumes of adamantylethylamine and mecamylamine occupy the voids during molecular stacking, displacing solvent molecules in the crystal structure of the smaller volume adamantyl Schiff base complexes. Additionally, the spatial site resistance attributed to the positional isomerism of adamantylethylamine and mecamtadine near the coordination site does not alter the coordination denticity between the metal and the ligand, but it does affect the coordination conformation of the Co(II) and Schiff base  $\kappa^1$ -O-monodentate complexes. In this context, the greater spatial site resistance of adamantylethylamine leads to a trans-coordination of the two Schiff base ligands, resulting in a distinct stacking pattern.

**Reversible solid-state conversion between  $\kappa^1$ -O-monodentate and  $\kappa^2$ -O, N-bidentate complexes.** The solid-state transformation was carried out by adding a proper ratio of base using complexes  $\text{CoCl}_2(\text{HL})_2$  (L=L1-L6) as the starting materials, respectively. As seen in Figure. 7, by grinding complexes  $\text{CoCl}_2(\text{HL})_2$  and the strong base NaOH in the molar ratio of 1:2 in the presence of two drop of methanol (60  $\mu\text{L}$ ), a color changed from green to red was visible within 5 minutes, indicating the generation of a new phase. PXRD analysis confirmed that the red sample obtained after grinding was the  $\kappa^2$ -O, N-bidentate complex  $\text{CoL}_2$ . The release of HCl was corroborated by the presence of NaCl ( $2\theta = 31.5^\circ$ ) which was formed as byproduct. In addition, the red  $\text{CoL}_2$  (L=L1-L6) solid, when ground with equimolar hydrochloric acid, undergoes a reverse reaction in the solid state, involving molecular rearrangement facilitated by the re-addition of HCl molecules to the structure, resulting in the production of the green complex  $\text{CoCl}_2(\text{HL})_2$ , which is further corroborated by PXRD (Figure S19). The application of mechanical force during the re-addition halogenation provides a novel synthetic route to the Schiff base cobalt  $\kappa^1$ -O-monodentate complexes, which cannot be synthesized through conventional solution methods. It is noteworthy that the direct grinding conversion of  $\text{CoCl}_2(\text{HL1})_2 \cdot \text{CH}_3\text{CN}$  to  $\text{CoCl}_2(\text{HL3})_2 \cdot \text{DMSO}$  could not be achieved due to the encapsulation of different solvent molecules in the solvated crystalline phase, and solid powder could not be obtained directly when utilizing DMSO as the LAG medium, attributed to its excellent solvation properties. Therefore, we employed both unsolvated



$\text{CoCl}_2(\text{HL1})_2$  and  $\text{CoCl}_2(\text{HL3})_2$  to modulate the dehalogenation process by disrupting the intramolecular hydrogen bonding, facilitating the transformation from  $\kappa^1\text{-O}$ -monodentate to  $\kappa^2\text{-O}$ , N-bidentate coordination. Other corresponding mono- and bi-dentate complexes can also undergo reversible dehalogenation and hydrogen conversion reactions, a process that has been confirmed by PXRD, as shown in Figures S20-S24.



**Figure 7.** Solid-state transformation from  $\text{CoCl}_2(\text{HL2})_2$  and  $\text{CoL2}$  by adding a proper ratio of base.

In the dehydrochlorination reaction, the potential intramolecular charge assisted  $\text{N-H}(+)\cdots\text{Cl}(-)$  interactions in  $\text{CoCl}_2(\text{HL})_2$  offer the opportunity to break hydrogen and coordination bonds and release H and Cl upon mechanochemical grinding with NaOH and then to form coordination bond of Co-N occurs to give the  $\text{CoL2}$ . Thus, with proper orientation and plausible  $\text{N-H}(+)\cdots\text{Cl}(-)$  interactions, it is assumed that the dehydrochlorination reaction that use charge assisted  $\text{N-H}(+)\cdots\text{Cl}(-)$  interactions can be applied to into precision synthesis of other similar Schiff base complexes. Under the influence of mechanical force, the addition of HCl leads to the protonation and subsequent breakdown of the Co-N coordination bond involving the nitrogen atom, which facilitates the formation of a new coordination bond between Cl and Co.



Additionally, the stability of the six-membered ring structure allows for the recovery of the N-H $\cdots$ O intramolecular hydrogen bonding interactions, resulting in the successful preparation of Schiff base  $\kappa^1$ -O-monodentate complexes.

### Antibacterial activity

Schiff base  $\kappa^1$ -O-monodentate and  $\kappa^2$ -O, N-bidentate complexes were used to test against one gram-positive (*Bacillus subtilis*) and one-gram negative bacteria (*Escherichia coli*) by the zone of inhibition method. The samples were prepared in concentrations of  $1.0 \times 10^{-1}$ ,  $1.0 \times 10^{-2}$ ,  $1.0 \times 10^{-3}$  and  $1.0 \times 10^{-4}$  mol L $^{-1}$  in DMF. The diameters of growth inhibition zones were measured after 48 h and the results were presented in Table S4 and Figure S43 and S44. Under the same conditions,  $\kappa^1$ -O-monodentate and  $\kappa^2$ -O, N-bidentate complexes showed better antibacterial effects against two bacteria compared with HL, which indicated that Schiff base ligand improved their biological activity after coordination with metal Co. Furthermore, the antibacterial effects of the two complexes were significantly concentration-dependent; as the concentration increased, the diameters of the bacterial inhibition zones also increased, demonstrating enhanced antibacterial activity. Notably, the  $\kappa^1$ -O-monodentate complexes **CoCl $_2$ (HL3) $_2$**  and **CoCl $_2$ (HL4) $_2$** , which contain memantine amine, exhibited the highest antimicrobial activity against *Escherichia coli*, with inhibition zone diameters of 18.0 mm and 20.0 mm, respectively, at a concentration of  $1.0 \times 10^{-1}$  mol L $^{-1}$  (Fig. 5). In contrast, the  $\kappa^2$ -O, N-bidentate complexes containing chlorosalicylaldehyde demonstrated superior antibacterial activity against *Bacillus subtilis*, with inhibition zone diameters of 24.0 mm for both **CoL1 $_2$**  and **CoL3 $_2$** .

### Conclusions

In this paper, we present a mechanochemical one-pot synthesis strategy that integrates the condensation covalent reaction of ammonia with aldehyde, metal coordination reactions, and dehydrohalogenation reactions. The raw materials employed include adamantylamine, 5-halosalicylaldehyde, and CoCl $_2 \cdot 6\text{H}_2\text{O}$ . Under the influence of mechanical forces, these reactions facilitate the self-ordering of the initial building blocks through synergistic interactions, leading to complex chemical processes that ultimately induce self-



classification and ultra-high selective coordination. This method successfully yielded 12 Co(II) complexes with Schiff bases, including the  $\kappa^1$ -O-monodentate complex **CoCl<sub>2</sub>(HL)<sub>2</sub>** and the  $\kappa^2$ -O, N-bidentate complex **CoL<sub>2</sub>**. Furthermore, this one-pot mechanochemical reaction not only simplifies the preparation procedure for Schiff base complexes but also enhances the complexity of mechanochemical multicomponent reactions. Notably, the reaction reached full conversion within 10 minutes, effectively demonstrating the advantages of mechanochemistry in green and sustainable synthesis. Additionally, the dormant chemical reactivity of the reactants was activated by mechanical force, resulting in the successful preparation of  $\kappa^1$ -O-monodentate complexes, which are typically challenging to obtain via solution methods. The reversible solid-state transformations between two complexes were studied by dehydrohalogenation/hydrohalogenation processes. The dehydrochlorination reaction from the complex **CoCl<sub>2</sub>(HL)<sub>2</sub>** to the complex **CoL<sub>2</sub>** involves the cleavage of the N–H and Co–Cl bonds of the complex **CoCl<sub>2</sub>(HL)<sub>2</sub>** and the formation of the coordination bonds Co–N of the **CoL<sub>2</sub>**. Meanwhile, the **CoL<sub>2</sub>** can also transform back to complex **CoCl<sub>2</sub>(HL)<sub>2</sub>** by the absorption of HCl via grinding. This work provides us a good example, showing reaction pathways in one-pot synthesis and solid-state transformation by understanding of mechanochemistry as a green route for the synthesis of Schiff base complexes.



## Experiment

### *Materials and Methods.*


All chemicals were obtained from commercial sources and used without further purification. Powder X-ray diffraction was performed with a Bruker D8 diffractometer ( $\lambda = 1.54056 \text{ \AA}$ ). Single crystal X-ray diffraction was carried out using a Bruker D8 QUEST X-ray single-crystal diffractometer. IR spectra were obtained with a PerkinElmer 100 FT-IR spectrometer using KBr pellets.

### *Mechanochemical synthesis of complexes $\text{CoCl}_2(\text{HL})_2$ (L = L1-L6).*

*“One-Pot” synthesis  $\text{CoCl}_2(\text{HL1})_2$ .* Amantadine (1mmol, 0.151 g), 5-chlorosalicylaldehyde (1mmol, 0.156 g) and  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  (0.5mmol, 0.119 g) in 2:2:1 molar ratio were ground in an agate mortar for 10 minutes, giving rise to green powder product, yield: 0.7105 g, 85%; M.p. 239.5-240.3 °C. IR (KBr): 3427 (m), 2906 (s), 2852 (m), 1643 (s), 1602 (w), 1526 (s), 1483 (s), 1324 (w), 1303 (m), 1229 (m), 1169 (s), 1112 (w), 1078 (m), 1032 (w), 833 (w), 793 (w), 682 (w), 627 (m), 521 (w), 492 (m), 446 (w). HRMS (ESI):m/z calculated for  $\text{C}_{34}\text{H}_{40}\text{Cl}_4\text{CoN}_2\text{O}_2 + \text{H}^+$ : 708.1176,  $[\text{M} + \text{H}^+]$  found: 708.1248.

*“One-Pot” synthesis  $\text{CoCl}_2(\text{HL2})_2$ .* Amantadine (1mmol, 0.151 g), 5-bromosalicylaldehyde (1mmol, 0.210 g) and  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  (0.5mmol, 0.119 g) in 2:2:1 molar ratio were ground in an agate mortar for 10 minutes, giving rise to green powder product, yield: 0.7105 g, 89%; M.p. 242.2-243.1 °C. IR (KBr): 3430 (m), 2048 (w), 2915 (s), 2853 (m), 2845 (m), 1641 (s), 1601 (m), 1529 (s), 1484 (s), 1454 (w), 1305 (m), 1233 (m), 1231 (m), 1171 (m), 1112 (m), 1079 (m), 940 (w), 834 (w), 794 (w), 683 (w), 628 (m), 523 (w), 495 (m), 447 (w). HRMS (ESI):m/z calculated for  $\text{C}_{34}\text{H}_{40}\text{Br}_2\text{Cl}_2\text{CoN}_2\text{O}_2 + \text{Na}$ : 783.5636,  $[\text{M} + \text{Na}]$  found: 783.0369

*“One-Pot” synthesis  $\text{CoCl}_2(\text{HL3})_2$ .* Memantine (1mmol, 0.216 g), 5-chlorosalicylaldehyde (1mmol, 0.156 g) and  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$  (0.5mmol, 0.119 g) in 2:2:1 molar ratio were ground in an agate mortar for 10 minutes, giving rise to green powder product, yield: 0.7105 g, 89%; M.p. 137.8-138.3 °C. IR (KBr): 3426 (m), 2917 (s), 2863 (w), 2845 (m), 2312 (w), 1643 (s), 1605 (s), 1531 (s), 1490 (s), 1455 (s), 1387 (s), 1301 (m), 1255 (w), 1231 (m), 1166 (s), 1024 (w), 937 (w),



870 (w), 825 (m), 799 (w), 735 (w), 644 (w), 585 (w), 541 (w). HRMS (ESI):m/z calculated for  $C_{38}H_{48}Cl_4CoN_2O_2+H$ : 764.1802,  $[M+H^+]$  found: 764.1874.

*“One-Pot” synthesis  $CoCl_2(HL4)_2$ .* Memantine (1mmol, 0.216 g), 5-bromosalicylaldehyde (1mmol, 0.2101 g) and  $CoCl_2 \cdot 6H_2O$  (0.5mmol, 0.119 g) in 2:2:1 molar ratio were ground in an agate mortar for 10 minutes, giving rise to green powder product, yield: 0.7105 g, 87%; M.p. 119.2-119.8 °C. IR (KBr): 3436 (s), 2944 (w), 2915 (s), 2863 (w), 2844 (w), 2300 (w), 1642 (s), 1602 (m), 1528 (s), 1486 (s), 1455 (s), 1386 (m), 1302 (s), 1255 (w), 1231 (m), 1165 (s), 1023 (w), 868 (w), 823 (m), 798 (w), 721 (w), 629 (w), 516 (w). HRMS (ESI):m/z calculated for  $C_{38}H_{48}Br_2Cl_2CoN_2O_2+Na$ : 874.0792,  $[M + Na]$  found: 874.0683

*“One-Pot” synthesis  $CoCl_2(HL5)_2$ .* Rimantadine (1mmol, 0.181 g), 5-chlorosalicylaldehyde (1mmol, 0.156 g) and  $CoCl_2 \cdot 6H_2O$  (0.5mmol, 0.119 g) in 2:2:1 molar ratio were ground in an agate mortar for 10 minutes, giving rise to green powder product, yield: 0.7105 g, 89%; M.p. 159.8-160.4 °C. IR (KBr): 3427 (m), 2905 (s), 2849 (m), 1647 (s), 1605 (w), 1529 (s), 1483 (s), 1386 (w), 1315 (m), 1283 (w), 1234 (m), 1173 (m), 1149 (w), 1089 (w), 1007 (s), 1034 (w), 871 (w), 826 (m), 798 (m), 714 (w), 689 (w), 637 (w), 547 (w). HRMS (ESI):m/z calculated for  $C_{38}H_{48}Cl_4CoN_2O_2+H$ : 764.1802,  $[M+H^+]$  found: 764.1877.

*“One-Pot” synthesis  $CoCl_2(HL6)_2$ .* Rimantadine (1mmol, 0.181 g), 5-bromosalicylaldehyde (1mmol, 0.210 g) and  $CoCl_2 \cdot 6H_2O$  (0.5mmol, 0.119 g) in 2:2:1 molar ratio were ground in an agate mortar for 10 minutes, giving rise to green powder product, yield: 0.7105 g, 89%; M.p. 167.9-168.4 °C. IR (KBr): 3444 (m), 2917 (s), 2863 (w), 2845 (m), 2312 (w), 1643 (s), 1605 (m), 1531 (s), 1489 (s), 1455 (s), 1388 (m), 1301 (m), 1255 (w), 1231 (m), 1166 (s), 1024 (w), 937 (w), 870 (w), 825 (m), 799 (w), 735 (w), 644 (w), 585 (w), 541 (w). HRMS (ESI):m/z calculated for  $C_{38}H_{48}Br_2Cl_2CoN_2O_2+Na$ : 874.0792,  $[M + Na]$  found: 874.0672

*Mechanochemical synthesis of complexes  $CoL_2$  ( $L = L1-L6$ ).*

*“One-Pot” synthesis  $CoL1_2$ :* Amantadine (1mmol, 0.1513 g), 5-chlorosalicylaldehyde (1mmol, 0.156 g),  $CoCl_2 \cdot 6H_2O$  (0.5mmol, 0.119 g) and NaOH (1mmol, 0.040 g) in 2:2:1:2 molar ratio were ground in an agate mortar aided by methanol (60 uL) for 5 minutes, yielding red powder products, yield: 0.6311 g, 88%, m.p. 290 °C. IR (KBr): 3434(m), 2908(s), 2850(m), 1598(s), 1522(s), 1455(s), 1389(s), 1347(w), 1315(m), 1268(w), 1243(w), 1166(m), 1132(w), 1107(m), 1077(m),





984(w), 934(w), 871(w), 823(m), 783(w), 753(m), 694(m), 662(m), 609(w), 548(m), 505(m), 452(w), 425(w). HRMS (ESI):m/z calculated for  $C_{34}H_{38}Cl_2CoN_2O_2+K$ : 674.2355, [M + K] found: 674.1273.

*“One-Pot” synthesis CoL2<sub>2</sub>*: Amantadine (1mmol, 0.151 g), 5-bromosalicylaldehyde (1mmol, 0.210 g),  $CoCl_2 \cdot 6H_2O$  (0.5mmol, 0.119 g) and NaOH (1mmol, 0.040 g) in 2:2:1:2 molar ratio were ground in an agate mortar aided by methanol (60 uL) for 5 minutes, yielding red powder products, yield: 0.6311 g, 89%, m.p. > 300 °C. IR (KBr): 3374(m), 3043(w), 2908(s), 2849(m), 1598(s), 1518(s), 1456(s), 1387(s), 1339(w), 1314(s), 1268(w), 1244(w), 1167(m), 1133(w), 1106(w), 1077(m), 985(w), 931(w), 872(w), 823(m), 781(w), 745(m), 685(w), 649(m), 605(w), 547(m), 499(m), 451(m), 424(w). HRMS (ESI):m/z calculated for  $C_{34}H_{38}Br_2CoN_2O_2+K$ : 762.1570, [M + K] found: 762.0263.

*“One-Pot” synthesis CoL3<sub>2</sub>*: Memantine (1mmol, 0.216 g), 5-chlorosalicylaldehyde (1mmol, 0.156 g),  $CoCl_2 \cdot 6H_2O$  (0.5mmol, 0.119 g) and NaOH (1mmol, 0.040 g) in 2:2:1:2 molar ratio were ground in an agate mortar aided by methanol (60 uL) for 5 minutes, yielding red powder products, yield: 0.6311 g, 86%, m.p. 236.7-238.4 °C. IR (KBr): 2909(m), 2863(w), 2846(m), 1601(s), 1523(s), 1456(s), 1411(w), 1387(s), 1316(m), 1297(w), 1254(w), 1166(s), 1132(w), 1093(w), 1055(w), 1026(w), 827(m), 809(w), 790(w), 734(w), 681(w), 663(w), 542(w), 514(w), 500(w). HRMS (ESI):m/z calculated for  $C_{38}H_{46}Cl_2CoN_2O_2+H$ : 692.2268, [M + H] found: 692.2341.

*“One-Pot” synthesis CoL4<sub>2</sub>*: Memantine (1mmol, 0.216 g), 5-bromosalicylaldehyde (1mmol, 0.210 g),  $CoCl_2 \cdot 6H_2O$  (0.5mmol, 0.119 g) and NaOH (1mmol, 0.040 g) in 2:2:1:2 molar ratio were ground in an agate mortar aided by methanol (60 uL) for 5 minutes, yielding red powder products, yield: 0.6311 g, 87%, m.p. 261.4-262.5 °C. IR (KBr): 2913(s), 1598(m), 1455(s), 1249(w), 1164(m), 1057(m), 1025(m), 985(w), 933(w), 824(m), 787(w), 648(w), 546(w), 505(w). HRMS (ESI):m/z calculated for  $C_{38}H_{46}Br_2CoN_2O_2+H$ : 780.1258, [M + H] found: 780.1331.

*“One-Pot” synthesis CoL5<sub>2</sub>*: Rimantadine (1mmol, 0.181 g), 5-chlorosalicylaldehyde (1mmol, 0.156 g),  $CoCl_2 \cdot 6H_2O$  (0.5mmol, 0.119 g) and NaOH (1mmol, 0.040 g) in 2:2:1:2 molar ratio were ground in an agate mortar aided by methanol (60 uL) for 5 minutes, yielding red powder products, yield: 0.6311 g, 86%, m.p. 258.3-260.7 °C. IR 3438(m), 2899(s), 2847(m), 1600(s), 1520(m), 1453(s), 1391(s), 1314(m), 1169(m), 1083(m), 979(w), 870(w), 815(w), 708(m), 657(w), 498(m),



456(m). HRMS (ESI):m/z calculated for  $C_{38}H_{46}Cl_2CoN_2O_2+H$ : 692.2268, [M + H] found: 692.2341.

*“One-Pot” synthesis CoL6<sub>2</sub>*: Rimantadine (1mmol, 0.181 g), 5-bromosalicylaldehyde (1mmol, 0.210 g),  $CoCl_2 \cdot 6H_2O$  (0.5mmol, 0.119 g) and NaOH (1mmol, 0.040 g) in 2:2:1:2 molar ratio were ground in an agate mortar aided by methanol (60 uL) for 5 minutes, yielding red powder products, yield: 0.6311 g, 89%, m.p. 264.3-265.6 °C. IR 3428(m), 2902(s), 2847(m), 1599(s), 1518(m), 1455(s), 1391(s), 1310(m), 1170(m), 1133(w), 1080(w), 975(w), 871(w), 824(w), 799(w), 684(m), 648(m), 496(m), 460(m). HRMS (ESI):m/z calculated for  $C_{38}H_{46}Br_2CoN_2O_2+H$ : 780.1258, [M + H] found:780.1331.

#### *Crystallization of $CoCl_2(HL)_2$ and $CoL_2$ (L = L1-L6).*

Recrystallization of six  $\kappa^1$ -O-monodentate complexes  $CoCl_2(HL)_2$  were carried out by evaporation of above green grinding product at ambient conditions using a variety of different solvents such as ethanol, acetone, dichloromethane, tetrahydrofuran, *N,N*-dimethylformamide, and dimethyl sulfoxide, and only in acetonitrile ( $CH_3CN$ ) gave rise to block-shape green single crystals suitable for single crystal X-ray (SC-XRD) analysis, in about 2 days.

Recrystallization of  $CoL_2$  was carried out by evaporation of the red grinding product in methanol/dimethyl sulfoxide (DMSO) (10:1) solution and gave rise to block-shaped red single crystals suitable for SC-XRD analysis, in about 7 days.

Recrystallization of  $CoL_3$ ,  $CoL_5$  and  $CoL_6$  were achieved through the evaporation of the red grinding product in an ethyl alcohol solution, resulting in the formation of block-shaped red single crystals suitable for single crystal X-ray diffraction (SC-XRD) analysis within approximately 5 days.

**Crystallography:** Single-crystal X-ray diffraction measurements of the crystals were performed with a Bruker Smart D8 Quest diffractometer equipped with a graphite monochromator. The determination of the unit-cell parameters and the data collection were performed with Mo- $K\alpha$  radiation ( $\lambda = 0.71073$  Å). The unit-cell parameters were obtained through least-squares refinements, and the structures were determined by direct methods and refined (on  $F^2$  with all independent data) by full-matrix least square methods (SHELXTL 2014). Data were reduced by



using the Bruker SAINT software. All the non-hydrogen atoms were directly located from different Fourier maps and refined with an isotropic displacement parameter. The hydrogen atoms in the ligand molecules were located in a difference electron density map and their positions were allowed to be refined together with individual isotropic temperature factors.

### *Antibacterial activity experiment*

Filter paper method was used to examine the antimicrobial activity against the two-gram positives (*Staphylococcus aureus* and *Bacillus subtilis*) and one-gram negative bacteria (*Escherichia coli*). The activated three bacteria were diluted to  $10^{-6}$  CFU/mL; 100  $\mu$ L of the bacterial solution and uniformly coated on the inactivated plate agar. Filter papers (6 mm) were immersed with HL,  $\text{CoCl}_2(\text{HL})_2$  and  $\text{CoL}_2$  or control group (cosolvent, *N,N*-dimethylformamide) were transferred to the Beef extract peptone AGAR medium agar plate. Thereafter, the agar plates were incubated at 37 °C for 48 h, the visible transparent halos on the plates were considered as the inhibition zones. The diameters and areas of the microbe-static ring were measured with Vernier calliper. Each sample was carried out in triplicates.

### ASSOCIATED CONTENT

Supporting Information. Further details on the synthesis and crystallographic information can be found in the Supporting Information. This material is available free of charge via the Internet at <http://pubs.acs.org>.”

### NOTES

The authors declare no competing financial interest.

### DATA AVAILABILITY



The data supporting this article have been included as part of the Supplementary Information. Crystallographic data for  $\text{CoCl}_2(\text{HL1})_2 \cdot \text{CH}_3\text{CN}$ ,  $\text{CoCl}_2(\text{HL2})_2 \cdot \text{CH}_3\text{CN}$ ,  $\text{CoCl}_2(\text{HL3})_2$ ,  $\text{CoCl}_2(\text{HL4})_2$ ,  $\text{CoCl}_2(\text{HL5})_2$ ,  $\text{CoCl}_2(\text{HL6})_2$ ,  $\text{CoL2}_2 \cdot \text{DMSO}$ , and  $\text{CoL3}_2$  have been deposited at the Cambridge Crystallographic Data Centre (CCDC) under the following accession numbers: 2083179, 2083171, 2343391, 2257072, 2121576, 2257071, 2074209, and 2343393. These data can be accessed via the CCDC website at [http://www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif). Additionally, the experimental crystal structure determinations for  $\text{CoL5}_2$  and  $\text{CoL6}_2$  are cataloged as CCDC experimental crystal structure determination: 969173 and 969172, respectively (C. Xu, L. Xiao-Chen, J. Xu-Dong, Y. Qi, H. Guang-Chao, G. Yu-Chen and H.-H. and Hu. Journal of Coordination Chemistry, 2014, 67, 352-362, DOI: 10.1080/00958972.2014.885509). These supplementary crystallographic data can be obtained free of charge from the CCDC.

## ACKNOWLEDGEMENTS

This research was supported by National Nature Science Foundation of China (No. 21571090), Liaoning BaiQianWan Talents Program, and Liaoning Provincial Department of Education Innovation Team Project (LT2017010), and the General project of the Liaoning Provincial Department of Education (LJKZ0102).

## AUTHOR INFORMATION

### *Corresponding Authors*

Daming Feng

ORCID: 0000-0001-7472-7278; E-mail: [dmfeng@lnu.edu.cn](mailto:dmfeng@lnu.edu.cn)

Xudong Jin

E-mail: [jinxudong@lnu.edu.cn](mailto:jinxudong@lnu.edu.cn)

Fang Guo

ORCID: 0000-0001-5014-2830; E-mail: [fguo@lnu.edu.cn](mailto:fguo@lnu.edu.cn)

### Author Contributions




The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

## REFERENCES

1. X. Liu, Y. Li, L. Zeng, X. Li, N. Chen, S. Bai, H. He, Q. Wang and C. Zhang, *Advanced Materials*, 2022, **34**, 2108327.
2. N. Fantozzi, J.-N. Volle, A. Porcheddu, D. Virieux, F. García and E. Colacino, *Chemical Society Reviews*, 2023, **52**, 6680-6714.
3. A. Krusenbaum, S. Grätz, G. T. Tigineh, L. Borchardt and J. G. Kim, *Chemical Society Reviews*, 2022, **51**, 2873-2905.
4. V. Martinez, T. Stolar, B. Karadeniz, I. Brekalo and K. Užarević, *Nature Reviews Chemistry*, 2023, **7**, 51-65.
5. D. Tan and F. García, *Chemical Society Reviews*, 2019, **48**, 2274-2292.
6. B. R. Naidu, T. Sruthi, R. Mitty and K. Venkateswarlu, *Green Chemistry*, 2023, **25**, 6120-6148.
7. L. E. Wenger and T. P. Hanusa, *Chemical Communications*, 2023, **59**, 14210-14222.
8. J. Zhang, L. Xu and W.-Y. Wong, *Coordination Chemistry Reviews*, 2018, **355**, 180-198.
9. M. Kumar, A. K. Singh, V. K. Singh, R. K. Yadav, A. P. Singh and S. Singh, *Coordination Chemistry Reviews*, 2024, **505**, 215663.
10. A. Abd-El-Aziz, Z. Li, X. Zhang, S. Elnagdy, M. S. Mansour, A. ElSherif, N. Ma and A. S. Abd-El-Aziz, *Topics in Current Chemistry*, 2025, **383**, 8.
11. I. Cocosila, A. Solé-Daura, P. Gotico, J. Forte, Y. Li and M. Fontecave, *ACS Catalysis*, 2024, **14**, 9618-9627.
12. S. Khan, X. Chen, A. Almahri, E. S. Allehyani, F. A. Alhumaydhi, M. M. Ibrahim and S. Ali, *Journal of Environmental Chemical Engineering*, 2021, **9**, 106381.
13. X. Liu and J.-R. Hamon, *Coordination Chemistry Reviews*, 2019, **389**, 94-118.
14. Y. Chen, H. Tang, H. Chen and H. Li, *Accounts of Chemical Research*, 2023, **56**, 2838-2850.
15. W. Ma, Y. Liu, N. Yu and K. Yan, *ACS Sustainable Chemistry & Engineering*, 2021, **9**, 16092-16102.



16. V. K. Singh, A. Chamberlain-Clay, H. C. Ong, F. León, G. Hum, M. Y. Par, P. Daley-Dee and F. García, *ACS Sustainable Chemistry & Engineering*, 2021, **9**, 1152–1160.
17. S. Zuo, S. Zheng, J. Liu and A. Zuo, *Beilstein Journal of Organic Chemistry*, 2022, **18**, 1860-5397.
18. Y. Gui, X. Yao, I. A. Guzei, M. M. Aristov, J. Yu and L. Yu, *Chemistry of Materials*, 2020, **32**, 7754-7765.
19. J.-F. Ayme and J.-M. Lehn, *Chemical Science*, 2020, **11**, 1114-1121.
20. L. Liang, W. Zhao, X.-J. Yang and B. Wu, *Accounts of Chemical Research*, 2022, **55**, 3218-3229.
21. J. Martí-Rujas and F. Guo, *Dalton Transactions*, 2021, **50**, 11665-11680.
22. S. Yang, Y. Yuan, X. Wang, Z. Hu and D. Guo, *Journal of Luminescence*, 2022, **242**, 118560.
23. A. A. Ahangar, A. A. Malik, I. Ahmad and A. A. Dar, *Dyes and Pigments*, 2023, **220**, 111742.
24. S. A. Boer, E. M. Foyle, C. M. Thomas and N. G. White, *Chemical Society Reviews*, 2019, **48**, 2596-2614.
25. G. Hubsher, M. Haider and M. S. Okun, *Neurology*, 2012, **78**, 1096-1099.
26. B. Kumar, K. Asha, M. Khanna, L. Ronsard, C. A. Meseko and M. Sanicas, *Archives of Virology*, 2018, **163**, 831-844.
27. H. D. Mai, N. M. Tran and H. Yoo, *Coordination Chemistry Reviews*, 2019, **387**, 180-198.
28. G. Peng, Y. Chen, B. Li, Y.-Q. Zhang and X.-M. Ren, *Dalton Transactions*, 2020, **49**, 5798-5802.
29. C. Xu, L. Xiao-Chen, J. Xu-Dong, Y. Qi, H. Guang-Chao, G. Yu-Chen and H.-H. and Hu, *Journal of Coordination Chemistry*, 2014, **67**, 352-362.
30. M. V. Novozhilova, E. A. Smirnova, J. A. Polozhentseva, J. A. Danilova, I. A. Chepurnaya, M. P. Karushev, V. V. Malev and A. M. Timonov, *Electrochimica Acta*, 2018, **282**, 105-115.
31. N. Singh, P. Kumar, S. Ahmad, J. Gupta, K. Raza and A. A. Hashmi, *Journal of Molecular Structure*, 2025, **1321**, 139874.
32. T. E. Shaw, J. Arami, J.-F. Ayme, J.-M. Lehn and T. Jurca, *RSC Mechanochemistry*, 2024, **1**, 33-37.





## DATA AVAILABILITY

View Article Online  
DOI: 10.1039/D5MR00057B

The data supporting this article have been included as part of the Supplementary Information. Crystallographic data for  $\text{CoCl}_2(\text{HL1})_2 \cdot \text{CH}_3\text{CN}$ ,  $\text{CoCl}_2(\text{HL2})_2 \cdot \text{CH}_3\text{CN}$ ,  $\text{CoCl}_2(\text{HL3})_2$ ,  $\text{CoCl}_2(\text{HL4})_2$ ,  $\text{CoCl}_2(\text{HL5})_2$ ,  $\text{CoCl}_2(\text{HL6})_2$ ,  $\text{CoL2}_2 \cdot \text{DMSO}$ , and  $\text{CoL3}_2$  have been deposited at the Cambridge Crystallographic Data Centre (CCDC) under the following accession numbers: 2083179, 2083171, 2343391, 2257072, 2121576, 2257071, 2074209, and 2343393. These data can be accessed via the CCDC website at [http://www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif). Additionally, the experimental crystal structure determinations for  $\text{CoL5}_2$  and  $\text{CoL6}_2$  are cataloged as CCDC experimental crystal structure determination: 969173 and 969172, respectively (C. Xu, L. Xiao-Chen, J. Xu-Dong, Y. Qi, H. Guang-Chao, G. Yu-Chen and H.-H. and Hu. Journal of Coordination Chemistry, 2014, 67, 352-362, DOI: 10.1080/00958972.2014.885509). These supplementary crystallographic data can be obtained free of charge from the CCDC.

